

University of São Paulo
School of Pharmaceutical Sciences of Ribeirão Preto
and
University of Technology Sydney
Centre for Forensic Science

Sweat as an alternative sample in doping control

Dayanne Cristiane Mozaner Bordin

2019

University of São Paulo
School of Pharmaceutical Sciences of Ribeirão Preto
and
University of Technology Sydney
Centre for Forensic Science

Sweat as an alternative sample in doping control

A Thesis presented to the Post Graduate
Program in toxicology and Forensic Sciences
in fulfilment of the requirements for the award
of the Doctor of Sciences and Doctor of
Philosophy

Area: Toxicology and Forensic Sciences

Supervisors: Prof. Dr. Bruno Spinosa De
Martinis and Dr. Philip Doble

Co- supervisors: Prof. Dr. Claude Roux

2019

ACKNOWLEDGEMENT

First and foremost, I would like to express my deepest appreciation to my supervisors, Professor Bruno Spinosa De Martinis and Dr. Lucas Blanes and my co-supervisors Professor Mauricio Yonamine and Professor Claude Roux. Thank you for giving me the opportunity to undertake this project, supporting and encouraging me throughout this journey. Thank you for giving me the freedom to try and follow my ideas and the opportunity to grow as a researcher.

I would like to thank Professor Philip Doble and Dr David Bishop for their constant laboratory support and invaluable contributions during the thesis preparation. I cannot put in words how grateful I am for everything you have done for me.

I also would like to show my gratitude to everyone involved in the preparation of this thesis. Your constructive feedback greatly improved the written work related to this PhD and I am extremely grateful for your time and efforts.

A sincere thank you to all the volunteers who participated in this study. During gathering this material, interviewing the participants, and collecting the data and samples, I had invaluable assistance from all of you. Without these, the completion of this study would not have been possible. I would like to thank Dr Claudio Sodré for the support with the laboratorial exams.

And to all my colleagues at the University of Sao Paulo (USP) and University of Technology Sydney (UTS) I cannot even begin to thank you all for everything you have done for me throughout my lifetime, but especially these past 4 years dealing with all my ups and downs. You are my “second family” you guys were always there for me made a world of difference.

I want to thank my partner Kevin for his support throughout the whole time. He has been by my side throughout the hardest year, to whom I can share my true emotions especially in the down times. His unconditional love, support, trust, and confidence in me and has helped and will continue to help me overcome many tough waves in life. I know the last few months have not been easy, and I greatly appreciate all the sacrifices you made to make my life easier even though they made your life harder. You have been there for me, always provided constructive feedback and advices.

Words cannot express how grateful I am and how much I love my mum Marcia, my dad Dorival, my sister Aryanne. It is because of them that today I have the chance to follow my dreams. I could not be where I am today without their love, hard work, support, and sacrifice.

I would like to thank all my family and friends back home in Brazil. Despite all these kilometres including an ocean separating us, you have been there for me every step of the way. You have always believed in me, even when I doubted myself.

“There are only two ways to live your life. One is as though nothing is a miracle. The other is as though everything is a miracle”.

Albert Einstein

ABSTRACT

Bordin, Dayanne Cristiane Mozaner. **Sweat as an alternative sample in doping control.** 2019. 149p. Thesis Doctor Degree. School of Pharmaceutical Sciences- USP and Centre of Forensic Sciences-UTS. 2019.

Sport has become a major industry of invaluable awards and major investments. To overcome opponents, many athletes resort to illegal means to improve their performance, as the World Antidoping Agency (WADA) prohibits drug use in sport. In addition to the ethical aspects, drug use can cause serious damage to an athletes' health by directly influencing physiological capacity and removing physical and psychological barriers, and their health damage may be irreversible. Urine is the standard matrix used in doping control; however, these analyses can also be performed on alternative biological samples; such as sweat. Toxicological analysis in sweat samples present some advantages over urine, such as less chance of sample tampering, greater detection windows, non-invasive collection, and the possibility of finding the parent drugs. The use of sweat as an alternative sample also allows obtaining a history of drug abuse.

The identification of doping cases has been an important area of research and development in recent years due to the continued increase in the number of prohibited substances. For the detection of prohibited substances, sophisticated analytical instruments such as liquid (LC), gas (GC) chromatography coupled to mass spectrometry and capillary electrophoresis (CE) are generally used. This thesis focused on the development of methods for evaluation of sweat as an alternative biological matrix for doping control, using different methods for sample preparation and detection to overcome its limitations. In addition, an outline of the profile of drugs used for sports practice and the acute effects caused in the organism of abusers were investigated.

A survey was applied to the volunteers to evaluate the adverse effects of androgenic anabolic steroids (AAS), dietary supplements (DS) and multiple drug use. In parallel, the volunteers hematological and biochemical parameters were measured. Through toxicological urine analysis, self-reporting and the parameters measured within the study revealed that the use of doping agents and drugs are subject to different factors, which are normally guided by the type of physical activity or aesthetic appearance. This abuse can increase the chances of health problems causing synergistic side effects, increasing the risk to develop diseases. We developed a method to determine the presence of 13 amphetamines and cocaine related

substances and their metabolites in sweat and urine using disposable pipette extraction (DPX) tips and GC-MS. The validated method was used to analyze 40 urine and sweat samples whose athletes self-reported the use of drugs and/or stimulants. It was verified that all consumed drugs and metabolites detected in urine were also present in sweat samples indicating that sweat was a viable matrix to perform doping tests.

We also developed a screening alternative method for stimulants detection in sweat samples using a microchip capillary electrophoresis instrument (Agilent Bionalyzer). Although functioning, the method was not sensitive enough to detect the low concentrations of drugs and metabolites present in sweat samples.

A one step fully automatized derivatisation and headspace (HS) SPME extraction method followed by GC-MS was developed for the analysis for amphetamine-type drugs and cocaine. The HS-SPME/GC-MS method was used to detect concentrations between 0.1 to 1 ng/mL of the target analytes without any additional sample preparation, suitable for routine analysis of drug traces in biological samples, such as urine and sweat.

Keywords: sport, doping, alternative biological matrix, sweat stimulants, urine, and analytical methods.

RESUMO

Bordin, Dayanne Cristiane Mozaner. **Suor como matriz alternativa no controle do doping**. 2019. 149p. Tese (Doutorado). Faculdade de Ciências Farmacêuticas de Ribeirão Preto – Universidade de São Paulo, Ribeirão Preto, 2019.

O esporte tornou-se uma indústria de valiosos prêmios e grandes investimentos. Na tentativa de vencer adversários, muitos atletas recorrem a meios ilegais para melhorar seu desempenho. A Agência Mundial Antidoping (WADA) proíbe o uso de drogas no esporte. Além dos aspectos éticos, o uso de drogas pode causar sérios danos à saúde de um atleta, influenciando diretamente a capacidade fisiológica e removendo barreiras físicas e psicológicas, e seus danos à saúde podem ser irreversíveis. A urina é a matriz padrão utilizada no controle de doping. No entanto, essas análises também podem ser realizadas em amostras biológicas alternativas; como o suor. A análise toxicológica em amostras de suor apresenta algumas vantagens em relação à urina, como menor chance de adulteração de amostras, maior janela de detecção, coleta não invasiva, além da possibilidade de encontrar principalmente as drogas consumidas. O uso de suor como amostra alternativa também permite obter um histórico de exposição ao abuso de drogas. A identificação de casos de doping tem sido uma área importante de pesquisa e desenvolvimento nos últimos anos devido ao aumento contínuo do número de substâncias proibidas. Para a detecção de substâncias proibidas, geralmente são utilizados instrumentos analíticos sofisticados, tais como cromatografia líquida (LC), cromatografia gasosa (GC) acoplados à espectrometria de massas e eletroforese capilar (CE). Esta tese centrou-se no desenvolvimento de métodos para avaliação do suor como uma matriz biológica alternativa para o controle de doping, utilizando diferentes métodos de preparação e detecção de amostras para superar suas limitações. Paralelamente, foi realizado um esboço do perfil das drogas utilizadas para prática esportiva e os efeitos agudos causados no organismo dos usuários. Foi aplicada uma pesquisa aos voluntários para avaliar os efeitos adversos dos esteroides anabolizantes androgênicos (AAS), suplementos alimentares (DS) e uso múltiplo de drogas. Em paralelo, foram realizadas medidas de parâmetros hematológicos e bioquímicos dos voluntários. Através de análises toxicológicas em urina, o auto-relato e a medição de parâmetros, o estudo revelou que o uso de agentes dopantes e drogas recreativas são submetidos a diferentes fatores, que são normalmente guiados pelo tipo de atividade física ou aparência estética. Seu uso pode aumentar as chances de problemas de saúde causando efeitos colaterais sinérgicos, aumentando o risco de desenvolver doenças. Desenvolvemos um

método para determinar a presença de 13 anfetaminas e substâncias relacionadas à cocaína e seus metabolitos em suor e urina usando Pipetas de Extração descartáveis (DPX) e GC-MS. O método validado foi utilizado para analisar 40 amostras de urina e suor cujos atletas auto-relataram o uso de drogas. Verificou-se que todas as drogas consumidas e seus metabólitos encontradas na urina também estavam presentes em amostras de suor indicando que o suor é uma matriz viável para realizar testes de doping. Também desenvolvemos um método alternativo de triagem para detecção de estimulantes em amostras de suor usando um instrumento de eletroforese capilar de microchip (Bionalyzer). Embora funcionasse, o método não era suficientemente sensível para detectar baixas concentrações de drogas e metabólitos presentes em amostras de suor. Foi desenvolvido um método de extração por headspace (HS) e derivatização totalmente automatizado de apenas um passo seguido por análises em GC-MS para determinação de drogas tipo anfetaminas e cocaína. O método HS-SPME/GC-MS foi utilizado para detectar concentrações entre 0,1 a 1 ng/mL dos analitos sem qualquer preparação adicional de amostra, apresentando potencial aplicação para análise de rotina de traços de drogas em amostras biológicas, como urina e suor.

Palavras-chave: esporte, doping, matriz biológica alternativa, estimulantes, agentes anabólicos esteroides, suor, urina e métodos analíticos.

CERTIFICATE OF AUTHORSHIP AND ORIGINALITY

I, Dayanne Cristiane Mozaner Bordin, declare that this thesis, is submitted in fulfilment of the requirements for the award of Doctor of Philosophy, in the Faculty of Science at the University of Technology Sydney.

This thesis has also been submitted for qualification at University of São Paulo as it is the result of a research candidature conducted as part of a collaborative Doctoral degree.

This document has not been submitted for qualifications at any other academic institution.

This thesis is wholly my own work unless otherwise reference or acknowledged. In addition, I certify that all information sources and literature used are indicated in the thesis.

This research is supported by an Australian Government Research Training Program, Brazilian Federal Agency for Coordination of Improvement of Higher Education Personnel (CAPES) and São Paulo Research Foundation (FAPESP).

Production Note:
Signature removed
prior to publication.

Dayanne Cristiane Mozaner Bordin

03.04.2019

LIST OF PUBLICATIONS

The presented work in this thesis have been published, accepted for publication or prepared for submission to journals as follows:

Published:

D.C.M. Bordin, F. Monedeiro, L.H.Bueno, M.N.R.Alves E.G. De Campos, B.S. De Martinis. Novas perspectivas em preparo de amostras biológicas com interesse forense, *Scientia Chromatographica*, 2015, DOI: **/10.4322/sc.2015.022**

D.C.M. Bordin, M.N.R.Alves, E.G. De Campos, B.S. De Martinis. Review Article: Disposable pipette tips extraction: Fundamentals, applications and state of the art, *Journal of Separation Science*, 2016, 39, 1168–1172, DOI: 10.1002/jssc.201500932

D.C.M. Bordin, B.B. Bettim, G.C. Perdoná, E.G. De Campos, B.S. De Martinis. Understanding alterations on blood and biochemical parameters in athletes that use dietary supplements, steroids and illicit drugs, *Toxicology*, 2017, **376**, 75-82, DOI:org/10.1016/j.tox.2016.05.019

A.A. Ishikawa, D.C.M. Bordin, E.G. De Campos, L. Blanes, P. Doble, B.S. De Martinis. Analysis of methylenedioxy derivatives in vitreous humor using liquid-liquid extraction and GC/MS. *Journal of Analytical Toxicology*, 2018, 42 (9), 661-666. Doi.org/10.1093/jat/bky044

Submitted and final preparation:

1. Dayanne Mozaner Bordin, Eduardo Geraldo de Campos, Lucas Blanes, Philip Doble, Claude Roux and Bruno Spinosa De Martinis. A new approach for determination of doping used-stimulants drugs in sweat and urine using pipette tips extraction (DPX) and GC/MS. (article) Drug Testing Analysis (**final preparation**)

2. Dayanne Mozaner Bordin, Eduardo Geraldo de Campos, Fernanda Monedeiro, Lucas Blanes, Philip Doble, Claude Roux and Bruno Spinoso De Martinis. Sweat as a tool for drug testing, metabolomics and diseases detection (**final preparation**)

3. Lidiane Alves, Dayanne Mozaner Bordin, Fernanda Monedeiro, and Bruno Spinoso De Martinis. Investigation of Ayahuasca β -carboline alkaloids and Tryptamines present in sweat samples after a religious ritual. Drug Testing Analysis (**final preparation**)

LIST OF CONFERENCES

The research conducted during this project was presented at several international conferences listed below.

Year	Conference	Presentation
2014	52 rd The International Association of Forensic Toxicologists (TIAFT), Buenos Aires, Argentina	A rapid assay for determination of MDMA, MDA, MDEA, Methamphetamine and Amphetamine in sweat using Disposable Pipette Extraction (DPX) and GC-MS
2015	53 rd The International Association of Forensic Toxicologists (TIAFT), Firenze, Italy	Side effects and trends in the consumption of anabolic-androgenic steroids (AASs) and psychotropic drugs in bodybuilders
2015	9th Congress of Toxicology in Developing Countries XIX Congresso Brasileiro de Toxicologia, Natal, Brazil	Side effects in the consumption of Anabolic Androgenic Steroids And Dietary Supplements
2016	54 rd The International Association of Forensic Toxicologists (TIAFT), Brisbane, Australia	A new approach for the analysis of stimulants in sweat using a disposable pipette extraction (DPX) and GC-MS
2016	International Symposium on Advances in Separation Science (ASASS), Hobart, Tasmania	Application of screening method for stimulants in sweat samples and urine using a modified technique DPX and GC-MS
2017	Forensic and Clinical Association Meeting (FACTA), Melbourne, Australia	One step automatized analysis of drug stimulants using HS-SPME/GC-MS

LIST OF FIGURES

Figure 1. 1: Percentage of substances identified by WADA in each drug class during the year 2014 (WADA, 2015).	8
Figure 1.2: Percentage of stimulants identified by WADA during the year 2014 (WADA, 2015).	10
Figure 1.3: Illustration of the general mechanism of action of AAS (Figure adapted from CUMMINGS, 2001).	12
Figure 1.4: Illustration of structures found on the skin; sweat glands: eccrine and apocrine; follicle and pores (Figure adapted from Mayo Foundation, 2007).	20
Figure 1.5: Photo of a PharmChek® adhesive used for the sweat collection.	24
Figure 1.6: Sweat collection procedure. (1) Clean the skin with isopropyl alcohol; (2 and 3) adhesive fastening; (4) adhesive removal; (5) adhesive storage (Figure modified from TAVARES, 2014).	25
Figure 1.7: The anti-doping workflow for sample analysis. The urine sample is divided in two aliquots (A and B). Sample A is used for the screening testing and the confirmatory analysis; while the sample B is kept for cross-examination in positive cases (WADA, 2015).	27
Figure 1. 8: Schematic diagram of the manual DPX extraction (BORDIN et al, 2016).	30
Figure 1.9: Schematic and modified diagram of the process of DPX extraction (BORDIN et al, 2016).	32
Figure 1.10: SPME extraction using the headspace mode (Figure adapted from, INDELICATO et al 2014).	33
Figure 1.11: Photos showing the upper part of the microchip used in the Bioanalysis equipment containing the reservoirs (1) and glass chip and their micro channels (2).	36
Figure 1.12: Schematic illustrating the micro-channels interconnected with the wells, injection and detection points (Figure taken from and adapted from the Agilent Technologies website, 2008).	37
Figure 1.13: Bioanalyzer Agilent ME. The electrodes are centred in the reservoirs of the microchip when closing the lid.	37
Figure 3.1: Self-reported motivations for use of AAS (n=20); several volunteers reported multiple motivations and “other motivation(s)”.	51

Figure 3.2: Total ion current chromatogram obtained after LLE extraction of a drug-free urine sample spiked with 100 ng/mL of: AMP; MAMP; MDMA; BE; THCCOOH and the deuterium internal standards AMP-d11, MDMA-d5, BE-d3 and THCCOOH-d3 (100ng/mL) for toxicological urine analysis.	56
Figure 3.3: Boxplot of total testosterone measures from non-users and abusers (AAS and DS) groups.	65
Figure 3.4: Correlation graph of creatinine and hematocrit variables.	67
Figure 5.1: Chromatographic separation of analytes in a sweat sample fortified with 100 ng/patch of all the analytes.....	82
Figure 6.1: Diagram of the analytical procedure for the extraction of simulants in sweat samples and the application in the Bioanalyzer microchip.	94
Figure 6.2: Electropherogram of running electrolyte solution using LED-IF (λ_{em} 470, λ_{em} 525). Conditions: 25 °C; injection time 2 seconds; injection voltage 1.5 kV; separation voltage 1.5 kV.....	95
Figure 6.3: Derivatisation reaction between FITC and methamphetamine.	96
Figure 6.4: Electropherogram of FITC solution using LED-IF (λ_{em} 470, λ_{em} 525), in two different runs. Conditions: running electrolyte (50 mM SDS plus 50 mM sodium tetraborate, pH 9.66); 25 °C; injection time 2 seconds; injection voltage 1.5 kV; separation voltage 1.5kV.....	96
Figure 6.5: Electropherogram of 50µg of derivatised AMP, MA, MDEA and MDMA using LED-IF (λ_{ex} 470, λ_{em} 525). Conditions: running electrolyte (50 mM SDS plus 50 mM sodium tetraborate, pH 9.66); 25 °C; injection time 2 seconds; injection voltage 1.5 kV; separation voltage 1.5 kV.....	97
Figure 7.1: Calibration curves obtained for all investigated analytes (AMP, MA, MDMA and COC) after the headspace SPME extraction and GC-MS analysis. ...	105
Figure 7.2: Comparison of the relative response obtained with the two different SPME coatings CAR-DVB-PDMS and PDMS (n=3) after the extraction procedure.	107
Figure 7.3: Metamphetamine derivatisation reaction.	108
Figure 7.4: Effect of the ethyl chloroformate (500 µg/mL) volume on the analytes response after PDMS-SPME extraction (n=3).....	109
Figure 7.5: Salting out effect in the concentrations of 2 and 3% of NaCl on the analytes response after PDMS-SPME extraction (n=3).....	110
Figure 7.6: Buffer addition effects on the analytes response after SPME extraction (n=3).....	111

Figure 7.7: Chromatographic separation obtained from a 500 ng/mL mixed standards injection (AMP, MA, MDMA, COC) after the derivatisation procedure and HS-SPME (PDMS) extraction. 112

LIST OF TABLES

Table 1.1: List of prohibited substances and methods published by WADA in 2016 (WADA, 2016).....	5
Table 1.2: Summary of main characteristics related to the different biological matrices (Table adapted from De MARTINIS, 2008).	17
Table 1.3: Summarizes important characteristics of sweat analysis, between them the type of collection device, application site of the patch on human body, time of wearing.....	21
Table 2.1: m/z ions selected to identify the target substances present in urine samples and the respective retention times.	47
Table 2.2: Self-report of 40 volunteers about lifetime consumption and current use of AAS, DS, pharmaceutical and recreational drugs.	49
Table 2.3: Self-report of 40 volunteers about daily use of dietary supplements (DS).	50
Table 2.4: Volunteers' self-reports of 20 athletes that consumed AAS.....	52
Table 2.5: Self-report of 20 volunteers who suffer adverse effects caused when using the combination of AAS and DS.....	53
Table 2.6: GC-MS validation method for the analysis of 5 different illicit drugs in urine.....	55
Table 3.1: Blood parameters measured in users and non-users in comparison with reference levels.....	62
Table 3.2: Analysis of uric acid, urea, creatinine and the glomerular filtration rate (GFR) in the blood of athletes that are users or non-users of AAS, DS and drugs of abuse.....	66
Table 3.3: Measurement results of liver enzymes alkaline phosphatase, gamma-glutamyl transferase, glutamico-oxalacetic transaminase and glutamic-pyruvic transaminase through the blood of athletes that are users or non-users of AAS, DS and drugs of abuse.....	69

Table 3.4: Measurement results of lipid levels in the blood of athletes that are users or non-users of AAS, DS and drugs of abuse.	70
Table 4.1: m/z of ions used to identify the analytes in urine and sweat samples and the respective retention times.	78
Table 4.2: Analytical validation results of the DPX-GC-MS method for the analysis of urine fortified with 13 stimulants.	83
Table 4.3: Analytical validation of the DPX-GC-MS method for the analysis of artificial sweat fortified with 13 stimulants.	84
Table 4.4: Analytical and reported results obtained from the volunteers.	86
Table 5.1: Chemical structures of the target compounds.	91
Table 5.2: Retention time of each analyte investigated.	93
Table 5.3: LODs and LOQs obtained with the Bioanalyzer.	98
Table 5.4: Drugs of abuse cut-off values proposed by SAMHSA.	99
Table 6.1: m/z of ions used to identify the analytes and the respective retention times.	103
Table 6.2: Analytical validation results of the HS-SPME/GC-MS method for the stimulants analysis.	112

SUMMARY

ACKNOWLEDGEMENT.....	iii
CERTIFICATE OF AUTHORSHIP AND ORIGINALITY	x
LIST OF PUBLICATIONS.....	xi
LIST OF CONFERENCES	1
Chapter 1: Introduction	2
1.1. Prohibited substances in the sport	4
1.2. Stimulants abuse in the sport	8
1.3. Anabolic androgenic steroids (AAS).....	11
1.4. Biological matrices analysed for doping control.....	14
1.4.1. Urine.....	14
1.4.2. Blood	15
1.5. Alternative Biological Matrices	16
1.5.1. Sweat as an alternative Biological Matrix	18
1.5.1.1. The Physiology of Sweat	18
1.5.1.2. Incorporation of Drugs into Sweat.....	20
1.5.1.3. Sweat Collection	23
1.5.1.4. Advantages and disadvantages of sweat as an alternative matrix.....	25
1.6. Techniques for analytical drug testing in biological matrices.....	26
1.6.1. Analytical workflows in doping control	26
1.6.2. Sample Preparation Methods	28
1.6.2.1. Liquid-liquid extraction (LLE)	28
1.6.2.2. Solid Phase Extraction (SPE).....	29
1.6.2.3. Disposable pipette tip devices.....	29
1.6.2.4. Solid Phase Micro Extraction (SPME)	32
1.6.3. Analytical methods applied in doping control	33

1.6.3.1. Gas Chromatography-mass spectrometry (GC-MS).....	34
1.6.3.2. Capillary Electrophoresis (CE) and Microchip Electrophoresis (ME)	35
1.6.3.2.1. Bioanalyzer Microchip Electrophoresis (ME).....	36
1.6.3.3. Liquid Chromatography-mass spectrometry (LC-MS).....	37
Research Aims	38
Ethical Aspects.....	40
CHAPTER 3: EVALUATION OF THE ADVERSE EFFECTS OF MULTIPLE DRUG USE USING A SELF-QUESTIONNAIRE AND TOXICOLOGICAL URINE ANALYSIS	
	41
Chapter 3: Introduction	42
3.1. Material and Methods.....	44
3.1.1. Chemicals and Reagents	44
3.1.2. Calibrators and quality controls	44
3.2. Individuals	44
3.2.1. Questionnaire and data collection	45
3.2.2. Urine samples collection.....	45
3.3. Development of a method for toxicological urine analysis	45
3.3.1. Urine Sample Preparation	45
3.4. Instrumentation.....	46
3.5. Data analysis.....	48
3.7. Results and Discussion.....	49
3.7.1. Questionnaire and data collection	49
3.7.1.1. Prevalence of substances use.....	49
3.7.1.2. Motivation for use of AAS	50
3.7.1.3. Volunteer's AAS abuse pattern	51
3.7.1.4. Effects and adverse effects of multi-drug use	53
3.7.2. Method Validation results	55
3.7.2.1. Toxicological urine analysis results using GC-MS.....	55
3.8. Conclusion.....	57

CHAPTER 4: VERIFICATION OF THE ALTERATIONS CAUSED BY MULTIPLE DRUG USE MEASURING HAEMATOLOGICAL AND BIOCHEMICAL PARAMETERS	58
Chapter 4: Introduction	59
4.1. Material and Methods.....	60
4.1.1. Individuals.....	60
4.1.2. Questionnaire and data collection	60
4.1.4. Statistics	61
4.2. Results and Discussion	61
4.2.1. Assessment of blood parameters and hormones levels	61
4.2.2. Biomarkers of renal function	66
4.2.3. Biomarkers of liver function	68
4.2.4. Blood lipid levels	70
4.3. Conclusion.....	71
CHAPTER 5: DETERMINATION OF DOPING USED-STIMULANTS IN SWEAT AND URINE USING PIPETTE TIPS EXTRACTION (DPX) AND GC- MS-EI.....	72
Chapter 4: Introduction	73
5.1. Material and Methods.....	73
5.1.1. Chemicals and Reagents.....	73
5.1.2. Calibrators and quality controls	74
5.2. Individuals	75
5.3. Sweat and Urine samples collection.....	75
5.4. Development of a method for toxicological analysis in sweat and urine samples.....	76
5.4.1. Sample Preparation	76
5.5. Instrumentation.....	77
5.6. Data analysis.....	79

5.7. Method validation.....	79
5.8. Results and Discussion	82
5.9. Conclusions	88
CHAPTER 6: THE APPLICATION OF LAB-ON-A-CHIP IN SWEAT SAMPLES FOR ANALYSIS OF PROHIBITED STIMULANTS IN SPORTS	89
Chapter 6: Introduction	90
6.1. Materials and Methods	91
6.1.1. Chemicals and reagents	91
6.1.2 Preparation of solutions and reagents.....	92
6.1.3 Standard solutions	92
6.1.4. Nile Blue solution.....	92
6.1.5. Electrolyte preparation	92
6.1.6 Fluorescein isothiocyanate I (FITC) stock solution	92
6.2. Instrumentation.....	93
6.2.1. Electrolyte solution analysis	95
6.2.2 Derivatisation	95
6.2.3. Solvent extractor selection	97
6.3.4. Limits of Detection (LOD) and Limit of Quantification (LOQ).....	98
CHAPTER 7: ONE STEP AUTOMATED SPME SAMPLE PREPARATION FOR STIMULANTS DRUGS ANALYSIS	100
Chapter 7: Introduction	101
7.1. Materials and Methods	102
7.1.1. Chemicals and Reagents.....	102
7.1.2. Solutions.....	102
7.2. Solid-Phase Micro Extraction procedure.....	102
7.3. Instrumentation.....	103

7.4. Data analysis.....	104
7.5. Method validation.....	104
7.6. Results and Discussion	106
7.6.1. Derivatisation.....	108
Chapter 8: Conclusions and Future Research	114
References	117
APPENDICES	131

ABBREVIATIONS

WADA	World Antidoping Agency
LC	Liquid Chromatography
GC	Gas Chromatography
MS	Mass Spectrometry
CE	Capillary Electrophoresis
DPX	Disposable Pipette Extraction Tips
AAS	Anabolic Androgenic Steroids
DS	Dietary Supplements
EI	Electro Ionization
HS	Headspace
SPME	Solid Phase Micro Extraction
IOC	International Olympic Committee
S	Substances
M	Methods
HIF	Hypoxia-inducible factor stabilizers
CG	Chorionic Gonadotrophin
GH	Growth Hormone
SERM	Selective estrogen receptor modulators
mL	Millilitre
CNS	Central Nervous System
DAT	Dopamine
NOR	Noradrenaline
SER	Serotonin
MA	Methamphetamine
MDA	3,4-methylenedioxyamphetamine

MDMA	3,4-methylenedioxymethamphetamine
MDEA	3,4-methylenedioxyethylamphetamine
BE	Benzoylecgonine
EME	Methylecgonine ester
DHEA	Dehydroepiandrosterone
cAMP	Cyclic adenosine monophosphate
RNA	Ribonucleic
LH	Luteinizing Hormone
FSH	Follicle-stimulating Hormone
T4	Thyroxine
T3	Triiodothyronine
IRMS	Isotope Ratio Mass Spectrometry
LLE	Liquid–Liquid Extraction□
SPE	Solid-Phase Extraction
RP	Reversed-Phase sorbent
CX	Cation Exchanger sorbent
WAX	Anion Exchanger sorbent
PDMS	Polydimethylsiloxane
PA	Polyacrylate
CAR	Carbowax
CI	Chemical Ionisation
QMS	Single Quadrupole Mass Spectrometer
S/N	Signal-to-noise
QQQ	Triple Quadrupole
ME	Microchip electrophoresis
EOF	Electroosmotic flow

MEKC	Micellar Electrokinetic Capillary Chromatography
UV	ultra-violet
CID	Collision-Induced Dissociation
PES	Performance Enhancing Substances
THCCOOH	11-nor-9-carboxy-9-tetrahydrocannabinol
d	Deuterium
mg	Milligram
C	Celsius
°	Degrees
MSTFA	N-methyl-N-(trimethylsilyl) trifluoroacetamide
µg	Microgram
IS	Internal Standard
M	Molar
rpm	Rotation per minute
v	Volume
mm	Millimetre
m	Metre
µm	Micrometre
SIM	Selected Ion-Monitoring
m/z	Mass-to-charge ratio
min	Minutes
LOD	Limit of Detection
LOQ	Limit of Quantification
PC	Positive Control
THC	Cannabis
BCAA	Amino acids

MCV	Mean corpuscular volume
MCH	Mean corpuscular haemoglobin
MCHC	Mean corpuscular haemoglobin concentration
GFR	Glomerular Filtration Rate
dL	Decilitre
L	Litre
CKD	Chronic kidney disease
GGT	Gama-Glutamyl Transferase
GOT	Glutamico-Oxalacetic Transaminase
GPT	Glutamic-Pyruvic Transaminase
HDL	High-density lipoprotein
LDL	Low-density lipoprotein
VLDL	Very Low-density lipoprotein
MPT	Mephentermine
AMP	Amphetamine
AMP D11	Amphetamine D11
METAMP	Metamphetamine
AMPHEP	Amphepramone
MDMA	3,4-Methylenedioxymethamphetamine
MDMA D5	3,4-Methylenedioxymethamphetamine D5
MDA	3,4-Methylendioxyamphetamine
MDEA	3,4-Methylenedioxyethylamphetamine
MPH	Methylphenidate
FENP	Fenproporex
COC	Cocaine
COC D3	Cocaine D3

BE	Benzoyllecgonine
COE	Cocaethylene
MEPHE	Mephedrone
SWGTOX	Scientific Working Group Toxicology guidelines
R ²	Coefficient of determination
QC	Quality control
ng	Nanogram
LOC	Lab-on-a-chip
mmol	Millimol
LED	Fluorescence detection
FITC	Fluoresceine isothiocyanate
mM	Millimol
kV	Kilovolts
ppt	Parts per trillion
cEF	ethyl chloroformate